

Finite time blow-up for radially symmetric solutions to a critical quasilinear Smoluchowski-Poisson system

Tomasz Cieřlak* & Philippe Laurençot†

October 19, 2008

Abstract

Finite time blow-up is shown to occur for radially symmetric solutions to a critical quasilinear Smoluchowski-Poisson system provided that the mass of the initial condition exceeds an explicit threshold. In the supercritical case, blow-up is shown to take place for any positive mass. The proof relies on a novel identity of virial type.

1 Introduction

We study the occurrence of blow-up in finite time for radially symmetric solutions to a generalized Smoluchowski-Poisson system

$$\partial_t u = \operatorname{div}(a(u) \nabla u - u \nabla v) \quad \text{in } (0, \infty) \times B(0, 1), \quad (1)$$

$$0 = \Delta v + u - M \quad \text{in } (0, \infty) \times B(0, 1), \quad (2)$$

$$\partial_\nu u = \partial_\nu v = 0 \quad \text{on } (0, \infty) \times \partial B(0, 1), \quad (3)$$

$$u(0) = u_0 \geq 0 \quad \text{in } B(0, 1), \quad \int_{B(0, 1)} v(t, x) \, dx = 0 \quad \text{for any } t \in (0, \infty), \quad (4)$$

where $B(0, 1)$ denotes the unit ball of $\mathbb{R}^n$, $n \geq 2$, and M the mean value of u_0 . The diffusion coefficient a belongs to $\mathcal{C}^2([0, \infty))$ and is assumed to be positive for simplicity (see Remark 5 below). The system (1)-(4) arises in astrophysics as a model of self-gravitating Langevin particles [1] and in biology [5] where it is also known as the parabolic-elliptic Keller-Segel chemotaxis model.

*Institute of Applied Mathematics and Mechanics, University of Warsaw, Banacha 2, 02-097 Warszawa, Poland. E-mail: T.Cieslak@impan.gov.pl

†Institut de Mathématiques de Toulouse, CNRS UMR 5219, Université de Toulouse, 118 route de Narbonne, F-31062 Toulouse Cedex 9, France. E-mail: Philippe.Laurencot@math.univ-toulouse.fr

A fundamental property of solutions to (1)-(4) is that

$$u(t) \geq 0, \quad \int_{B(0,1)} u(t, x) \, dx = M |B(0,1)|, \quad \text{and} \quad \int_{B(0,1)} v(t, x) \, dx = 0 \quad (5)$$

for $t > 0$, which readily follows from (1), (3), the comparison principle, the non-negativity of u_0 , and the definition of v . It is by now well-known that, if $a \equiv 1$, there are radially symmetric initial data u_0 for which the corresponding solution to (1)-(4) blows up in finite time and this singular phenomenon may happen for any $M > 0$ if $n \geq 3$ but only for $M > 8\pi$ if $n = 2$ [4, 6]. More recently, it was shown in [3] that there is a critical exponent for the nonlinear diffusion coefficient a which separates two different behaviours for the solutions to (1)-(4): on the one hand, if $a(z) \geq c(1+z)^\alpha$ and $\alpha > (n-2)/n$, there is a unique global classical solution to (1)-(4) for any non-negative initial condition $u_0 \in L^\infty(B(0,1))$ (and this is actually true for a general smooth bounded domain of $\mathbb{R}^n$, $n \geq 1$). On the other hand, if $a(z) \leq c(1+z)^\alpha$ and $\alpha < (n-2)/n$, radially symmetric solutions to (1)-(4) blowing up in finite time are constructed in [3].

Except for $n = 2$, the critical case $\alpha = (n-2)/n$ is not covered in [3] and the purpose of this note is to fill this gap: indeed, the main outcome of our analysis is that, if $a(z) \leq c(1+z)^{(n-2)/n}$, there are solutions to (1)-(4) blowing up in finite time when M exceeds an explicit threshold. As a by-product of our analysis, we also establish an alternative and simpler proof of the blow-up result in [3] for $\alpha \in [0, (n-2)/n)$. Indeed, the construction of solutions to (1)-(4) blowing up in finite time performed in [3] relies on the possibility of reducing (1)-(4) to a single parabolic equation. The approach used in this paper is completely different and relies on the derivation of a differential inequality of virial type which cannot hold true for all times. When $a \equiv 1$, this technique is used in [6] where it is shown that the moment of order n of u satisfies a differential inequality which contradicts the non-negativity of u after a finite time. Seemingly, the moment of order n of u does not give valuable information when the diffusion is nonlinear and we introduce nonlinear functions of u to be able to handle this case. We finally point out that the above results are only valid for $n \geq 2$: the situation is qualitatively different in the one dimensional case $n = 1$ and will be considered in a separate paper [2].

2 Finite time blow-up

We first introduce some notation: for $(\alpha, p) \in [0, \infty) \times (1, \infty)$, we define

$$\kappa_p(\alpha) := \frac{(p-1)}{(\alpha+1)(p+\alpha)} \left(\frac{2(n-1)}{p-1} \right)^{\alpha+1} (np)^{(n-2-\alpha n)/n}. \quad (6)$$

We also define $\bar{m}_p(f)$ for $p \geq 1$ and $f \in L^\infty(0,1)$ by

$$\bar{m}_p(f) := \frac{1}{p} \int_0^1 \left(\int_r^1 f(\rho) \rho^{n-1} \, d\rho \right)^p r^{n-1} \, dr. \quad (7)$$

Our main result then reads as follows.

Theorem 1 *Assume that there are $\alpha \in [0, (n-2)/n]$, and positive real numbers $c_1 > 0$ and $c_2 > 0$ such that*

$$0 < a(z) \leq c_1 z^\alpha + c_2 \quad \text{for } z \geq 0. \quad (8)$$

Let $M > 0$ and consider a non-negative radially symmetric function $u_0 \in L^\infty(B(0, 1))$ such that $\|u_0\|_1 = M$. Assume further that $E_{M,p}(\bar{m}_p(u_0)) < 0$ for some $p > 1$, where

$$\begin{aligned} E_{M,p}(z) &:= c_1 \kappa_p(\alpha) \left(\frac{M}{n}\right)^{(2p+n\alpha(p+1))/n} z^{(n-2-\alpha n)/n} + c_2 \kappa_p(0) \left(\frac{M}{n}\right)^{(2p)/n} z^{(n-2)/n} \\ &+ M z - \frac{1}{p(p+1)} \left(\frac{M}{n}\right)^{p+1}, \quad z \geq 0. \end{aligned} \quad (9)$$

Then the system (1)-(4) has a unique maximal classical solution (u, v) with finite maximal existence time $T_{max} \in (0, \infty)$ and $\|u(t)\|_\infty \rightarrow \infty$ as $t \rightarrow T_{max}$.

Remark 2 *There are initial data u_0 for which $E_{M,p}(\bar{m}_p(u_0)) < 0$ for some $p > 1$. Indeed, observe that $E_{M,p}(0) < 0$ for all $M > 0$ if $\alpha \in [0, (n-2)/n]$ and for $M > n(c_1 p(p+1)\kappa_p((n-2)/n))^{n/2}$ if $\alpha = (n-2)/n$. It is then sufficient to take u_0 sufficiently concentrated near $x = 0$ so that $\bar{m}_p(u_0)$ is close to zero (for instance, $u_0 = M\delta^{-n} \mathbf{1}_{B(0,\delta)}$ for $\delta > 0$ sufficiently small).*

Proof of Theorem 1.

By [3, Theorem 1.3], there exist a maximal existence time $T_{max} \in (0, \infty]$ and a unique radially symmetric classical solution $(u, v) \in \mathcal{C}([0, T_{max}); L^2(B(0, 1); \mathbb{R}^2)) \cap \mathcal{C}^{1,2}((0, T_{max}) \times B(0, 1); \mathbb{R}^2)$ to (1)-(4) satisfying (5) for $t \in [0, T_{max})$. Moreover, if $T_{max} < \infty$ then $\|u(t)\|_\infty \rightarrow \infty$ as $t \rightarrow T_{max}$.

We introduce

$$U(t, r) := \frac{1}{n|B(0, 1)|} \int_{B(0,r)} u(t, x) \, dx \quad \text{and} \quad m_p(t) := \frac{1}{p} \int_0^1 \left(\frac{M}{n} - U(t, r)\right)^p r^{n-1} \, dr$$

for $(t, r) \in [0, T_{max}) \times [0, 1]$ and derive the following identity of virial type.

Lemma 3 *Let A be defined by $A' = a$ and $A(0) = 0$. Then*

$$\frac{dm_p}{dt} = M m_p - \frac{1}{p(p+1)} \left(\frac{M}{n}\right)^{p+1} + \mathcal{R}_p(u) \quad (10)$$

with

$$\mathcal{R}_p(u) := \int_0^1 r^{2n-3} \left(\frac{M}{n} - U\right)^{p-2} \left(2(n-1) \left(\frac{M}{n} - U\right) - (p-1) r^n u\right) A(u) \, dr.$$

Proof. Integrating (1) gives that U solves

$$\partial_t U = r^{n-1} \partial_r A(u) + u U - \frac{M}{n} r^n u \quad \text{with} \quad U(t, 0) = \frac{M}{n} - U(t, 1) = 0.$$

Consequently,

$$\begin{aligned} \frac{dm_p}{dt} &= \int_0^1 \left(\frac{M}{n} - U \right)^{p-1} \left[\frac{M}{n} (r^n - 1) \partial_r U + \left(\frac{M}{n} - U \right) \partial_r U - r^{2(n-1)} \partial_r A(u) \right] dr \\ &= -\frac{1}{p} \left(\frac{M}{n} \right)^{p+1} + M m_p + \frac{1}{p+1} \left(\frac{M}{n} \right)^{p+1} \\ &\quad + \int_0^1 \left[2(n-1) r^{2n-3} \left(\frac{M}{n} - U \right)^{p-1} - (p-1) r^{2(n-1)} \left(\frac{M}{n} - U \right)^{p-2} \partial_r U \right] A(u) dr, \end{aligned}$$

and hence (10). $\square$

The next step is to estimate $\mathcal{R}_p(u)$ in terms of m_p . To this end, we notice that the assumption (8) on a warrants that

$$0 \leq A(z) \leq \frac{c_1}{1+\alpha} z^{1+\alpha} + c_2 z, \quad z \geq 0. \quad (11)$$

In view of (11), we have

$$\begin{aligned} &\left[2(n-1) \left(\frac{M}{n} - U \right) - (p-1) r^n u \right] A(u) \\ &\leq \max \left\{ 2(n-1) \left(\frac{M}{n} - U \right) - (p-1) r^n u, 0 \right\} \left(\frac{c_1}{1+\alpha} u^{1+\alpha} + c_2 u \right) \\ &\leq \max \left\{ 2(n-1) \left(\frac{M}{n} - U \right) - (p-1) r^n u, 0 \right\} \left(\frac{c_1}{1+\alpha} \left[\frac{2(n-1)}{(p-1)r^n} \left(\frac{M}{n} - U \right) \right]^\alpha + c_2 \right) u \\ &\leq \frac{c_1(p-1)}{1+\alpha} \left(\frac{2(n-1)}{p-1} \right)^{1+\alpha} \left(\frac{M}{n} - U \right)^{1+\alpha} r^{-n\alpha} u + 2(n-1)c_2 \left(\frac{M}{n} - U \right) u, \end{aligned}$$

and thus

$$\begin{aligned} \mathcal{R}_p(u) &\leq \frac{c_1(p-1)}{1+\alpha} \left(\frac{2(n-1)}{p-1} \right)^{1+\alpha} \int_0^1 r^{n-2-\alpha n} \left(\frac{M}{n} - U \right)^{p+\alpha-1} \partial_r U dr \\ &\quad + 2(n-1)c_2 \int_0^1 r^{n-2} \left(\frac{M}{n} - U \right)^{p-1} \partial_r U dr. \end{aligned} \quad (12)$$

Since $\alpha \in [0, (n-2)/n]$, the function $r \mapsto r^{(n-2-\alpha n)/n}$ is concave and we infer from the Jensen inequality

(with measure $((M/n) - U)^{p+\alpha-1} \partial_r U \, dr$) that

$$\begin{aligned}
& \int_0^1 r^{n-2-\alpha n} \left(\frac{M}{n} - U \right)^{p+\alpha-1} \partial_r U \, dr \\
& \leq \left(\frac{1}{p+\alpha} \left(\frac{M}{n} \right)^{p+\alpha} \right)^{(2+\alpha n)/n} \left(\int_0^1 r^n \left(\frac{M}{n} - U \right)^{p+\alpha-1} \partial_r U \, dr \right)^{(n-2-\alpha n)/n} \\
& \leq \left(\frac{1}{p+\alpha} \left(\frac{M}{n} \right)^{p+\alpha} \right)^{(2+\alpha n)/n} \left(\frac{n}{p+\alpha} \int_0^1 r^{n-1} \left(\frac{M}{n} - U \right)^{p+\alpha} \, dr \right)^{(n-2-\alpha n)/n} \\
& \leq \frac{(np)^{(n-2-\alpha n)/n}}{p+\alpha} \left(\frac{M}{n} \right)^{(2p+\alpha n(p+1))/n} m_p^{(n-2-\alpha n)/n}.
\end{aligned}$$

Arguing in a similar way to estimate the second integral in the right-hand side of (12), we deduce from (12) that

$$\mathcal{R}_p(u) \leq E_{M,p}(m_p) - M \, m_p + \frac{1}{p(p+1)} \left(\frac{M}{n} \right)^{p+1}.$$

Inserting this estimate in (10) we arrive at

$$\frac{dm_p(t)}{dt} \leq E_{M,p}(m_p(t)) \quad \text{for } t \in [0, T_{max}). \quad (13)$$

Assume now for contradiction that $T_{max} = \infty$. Since $z \mapsto E_{M,p}(z)$ is an increasing function and $m_p(0) = \bar{m}_p(u_0)$, we realize that, as soon as $E_{M,p}(m_p(0)) < 0$, we have $m_p(t_0) = 0$ for some $t_0 \in (0, \infty)$. Thus, $U(t_0, r) = M/n$ for all $r \in [0, 1]$ which contradicts the fact that $U(t_0, 0) = 0$. Consequently, $T_{max} < \infty$ and the proof is complete. $\square$

Remark 4 If $a \equiv 1$ (i.e. $\alpha = 0$), Lemma 3 is also valid for $p = 1$ and m_1 coincides with the moment of order n used in [6]. Not surprisingly, if $n = 2$ and $a \equiv 1$, we have $E_{M,p}(0) < 0$ if $M > 4(p+1)$ which converges to 8 as p approaches 1 and we recover the well-known threshold condition $\|u_0\|_1 > 8\pi$ for finite time blow-up to occur in the parabolic-elliptic Keller-Segel system [6].

Remark 5 The requirement $a > 0$ is only used to have classical solutions to (1)-(4) but does not play any role in the blow-up condition and the identity of virial type (Lemma 3). Thus, Theorem 1 remains valid if the diffusion is degenerate (for instance, $a(r) = mr^{m-1}$ with $m \in [1, 2(n-1)/n]$) provided an appropriate notion of weak solutions is available.

Acknowledgement. This paper was prepared during T. Cieřlak's one-month visit at the Institut de Mathématiques de Toulouse, Université Paul Sabatier. T. Cieřlak would like to express his gratitude for the invitation, support, and hospitality.

References

- [1] P.-H. Chavanis, *Generalized thermodynamics and Fokker-Planck equations. Applications to stellar dynamics and two-dimensional turbulence*, Phys. Rev. E **68** (2003), 036108.
- [2] T. Cieřlak and Ph. Laurençot, in preparation.
- [3] T. Cieřlak and M. Winkler, *Finite-time blow-up in a quasilinear system of chemotaxis*, Nonlinearity **21** (2008), 1057–1076.
- [4] W. Jäger and S. Luckhaus, *On explosions of solutions to a system of partial differential equations modelling chemotaxis*, Trans. Amer. Math. Soc. **329** (1992), 819–824.
- [5] E.F. Keller and L.A. Segel, *Initiation of slime mold aggregation viewed as an instability*, J. Theor. Biol. **26** (1970), 399–415.
- [6] T. Nagai, *Blow-up of radially symmetric solutions to a chemotaxis system*, Adv. Math. Sci. Appl. **5** (1995), 581–601.